



URETERAL STONES

Farzin Soleimanzadeh MD

Tabriz University of Medical Sciences

Diagnosis

- Solitary kidney / Anuria
- Fever

Diagnosis

- Solitary kidney / Anuria
- Fever
- U/A or U/C?

Diagnosis


- Solitary kidney / Anuria
- Fever
- U/A or U/C?
- Imaging modalities:
 - Sono (first): sens. 45, spec. 94 for ureteral stones
sens. 45, spec. 88 for renal stones
 - KUB: sens. 44, spec. 77 for ureteral stones (only for opacity and follow up IF CT available)
 - NCCT: sens. 93.1, spec. 96.6 for ureteral stones, sens. 100% for calculi > 3 mm
 - X-ray concern: Low dose

Management of Renal Colic




- NSAIDS
 - (**caution:** Coronary events, **contraindication:** Diclofenac in IHD, CHF, CVD, vascular disease)
- Paracetamol (Acetaminophen)

Management of Renal Colic




- NSAIDS
 - (**caution:** Coronary events, **contraindication:** Diclofenac in IHD, CHF, CVD, vascular disease)
- Paracetamol (Acetaminophen)
- Opioids
- Ketamine + Opioids
- Ketamine + NSAIDS

Management of Renal Colic



- NSAIDS
 - (**caution:** Coronary events, **contraindication:** Diclofenac in IHD, CHF, CVD, vascular disease)
- Paracetamol (Acetaminophen)
- Opioids
- Ketamine + Opioids
- Ketamine + NSAIDS
- Antispasmodics: NO
- Lidocaine IV injection??

Management of Renal Colic



- NSAIDS
 - (**caution:** Coronary events, **contraindication:** Diclofenac in IHD, CHF, CVD, vascular disease)
- Paracetamol (Acetaminophen)
- Opioids
- Ketamine + Opioids
- Ketamine + NSAIDS
- Antispasmodics: NO
- Lidocaine IV injection??
- Acupuncture?

Preventing Recurrent Renal Colic

- NSAIDS (pills / supp)
 - Diclofenac and renal function

Preventing Recurrent Renal Colic



- NSAIDS (pills / supp)
 - Diclofenac and renal function
- MET and pain recurrence prevention

Preventing Recurrent Renal Colic

- NSAIDS (pills / supp)
 - Diclofenac and renal function
- MET and pain recurrence prevention
- Resistant pain: no recovery with ORAL medication:
 - Drainage (Stenting / PCN)
 - Stone removal

Concomitant UTI

- Stent or PCN?
- Timing of STONE removal



Medical Expulsive Therapy

- Only in informed patients
- α -blockers
- calcium-channel inhibitors (nifedipine)
- phosphodiesterase type 5 inhibitors (PDEI-5) (tadalafil)

Medical Expulsive Therapy

- Only in informed patients
- α -blockers
- calcium-channel inhibitors (nifedipine)
- phosphodiesterase type 5 inhibitors (PDEI-5) (tadalafil)
- Steroids?

Medical Expulsive Therapy

- Only in informed patients
- α -blockers
- calcium-channel inhibitors (nifedipine)
- phosphodiesterase type 5 inhibitors (PDEI-5) (tadalafil)
- Steroids?

Recommendation	Strength rating
Offer α -blockers as medical expulsive therapy as one of the treatment options for (distal) ureteral stones > 5 mm.	Strong

Best Practice in Ureteroscopy

- Rigid: size, location of stone
- Flexible

Best Practice in Ureteroscopy

- Rigid: size, location of stone
- Flexible
- Fluoroscopic Equipment
- Safety guide wire

Best Practice in Ureteroscopy

- Rigid: size, location of stone
- Flexible
- Fluoroscopic Equipment
- Safety guide wire
- Antegrade ureteroscopy / laparoscopy (in upper, > 15 mm, impaction)
- Stenting and retrieval
- Intracorporeal lithotripsy (Holmium YAG laser versus pneumatic and US)
- Stenting (ureteral trauma, residual fragments, bleeding, perforation, UTIs, or pregnancy)

Complications of URS

- Overall: 9 – 25%
- Mostly minor, not necessitating intervention
- Ureteral avulsion and strictures are rare (< 1%)
- Previous perforations are the most important risk factor for complications

What we still don't know

- How long to wait with MET


What we still don't know

- How long to wait with MET
- How large is large?

What we still don't know

- How long to wait with MET
- How large is large?
- SWL Vs URS? (regarding overall stone free rate, faster results, complications, obese and morbid obese)

Proximal Ureteral Stone



Distal Ureteral Stone



Thank you

